

## DATA SHEET

Version: 03  
Revision date: 15/06/2023

Tlf: +34 983 54 85 63  
[info@canvaxbiotech.com](mailto:info@canvaxbiotech.com)

[www.canvaxbiotech.com](http://www.canvaxbiotech.com)


### 1. Identification

**Product name** **pSpark® IV**  
(20ng/μL) 20 reactions

**Cat No** **C0004**

### 2. Description

**pSpark® IV** is a highly efficient, accurate and easy-to-use DNA cloning system that exploits its very low background feature for the expression of toxic genes under transcription-free conditions. In this vector, the lac promoter has been eliminated and therefore blue/white screening is not allowed (alpha-peptide coding region remains and you can find blue colony). The vector is ideal for cloning genes that produce toxic polypeptides by transcription/translation.



### 3. Composition

Item	Quantity
pSpark® IV DNA cloning vector	20 rxn (20ng/μL)
10X PEG 6000 solution	150 μL
1kb control insert	5 μL
T4 DNA ligase (5 Weiss U/μL)	20 μL
10x T4 DNA ligase Buffer	100 μL

### 4. Features

- **Unprecedented high cloning efficiency:** > 2,500 positive colonies expected under optimal conditions.
- **Transcription-free.**
- **Easy-to-use:** eliminate screening of recombinants due to its
- **High stability:** eliminates cloning bias or pitfalls.
- **Time-saving protocol:** avoids any step required after PCR, just 19 minutes from PCR to plating.
- **Powerful:** obtain 5x more positive colonies using 10x less DNA insert.
- **Great versatility:** compatible with any protocol, proofreading polymerase, competent cells, ligation time or primers.
- **Sensitive:** clone from 50 bp insert to up to 14 kb with just 5ng per kb of insert.
- **Cost avoidance:** removes expensive primer phosphorylation use.
- **Eliminates positive selection vector.**



## DATA SHEET

Version: 03

Revision date: 15/06/2023

Tlf: +34 983 54 85 63  
[info@canvaxbiotech.com](mailto:info@canvaxbiotech.com)[www.canvaxbiotech.com](http://www.canvaxbiotech.com)

### 5. Storage specifications

Store in a non-frost-free freezer at **-20°C**.

### 6. Applications

- Cloning of High-Fidelity PCR amplified products.
- Production of ssDNA.
- *In vitro* transcription from T7/SP6 dual-opposed promoters.
- Cloning of toxic genes.

### 7. Recommended protocol

#### 1. Amplification

Amplify DNA with any proofreading polymerase following suggested protocol. Gel purifies the insert or check by electrophoresis the quality of amplified DNA. If a single band appears on the gel and template DNA is not a vector with Ampicillin resistance, like the pSpark® IV DNA cloning vector, then gel purification is not needed.

#### 2. Ligation

Suggested ratio of insert to vector is 5:1. Use the equation **ng insert = 33ng x kb** (of insert, that is, for a 1kb insert only 33 ng are needed) or use **1 µL** of unpurified PCR product for directly ligation.

##### Mix:

**1 µL** of pSpark® IV DNA cloning vector (20ng/µL)

**1 µL** 10x T4 DNA ligase buffer

**1 µL** 10x PEG 6000 solution

**X µL** of insert

**1 µL** of T4 DNA ligase (5 Weiss units)

H<sub>2</sub>O to **10 µL**

Incubate the reaction 60 minutes at 22°C (a ligation time from 10min to overnight could be used)

#### 3. Transformation

Transformation can be made with your regular protocol, competent cells and selective media. As pSpark® IV DNA cloning vector has Ampicillin resistance, this antibiotic is needed. Also, as pSpark® IV DNA cloning vector has blue/white screening capability, IPTG/XGal plates are recommended.

A suitable transformation protocol is:

- Thaw competent cells on ice, take **50 µL** and put into a cool 1,5 mL tube.
- Add **10 µL of the ligation mixture**, mix by gently flicking with fingertips and incubate on ice for 30 minutes.
- Heat-shock the cells at **exactly 42°C for exactly 45 seconds** (do not shake) and then transfer to ice for 2 minutes.
- Plate the cells into media with antibiotic, IPTG and XGal and incubate overnight at 37°C.

#### 4. Analysis of transformants

Either colony PCR or plasmid purification and digestion could be used for analysis of transformants. pSpark® IV DNA cloning vector has pUC/M13 forward and reverse sequencing primers binding site for sequencing or amplification of insert. Insert size when amplified with pUC/M13 fw and rev primers is about 200bp longer than the real size of insert.



## DATA SHEET

Version: 03

Revision date: 15/06/2023

Tlf: +34 983 54 85 63  
[info@canvaxbiotech.com](mailto:info@canvaxbiotech.com)[www.canvaxbiotech.com](http://www.canvaxbiotech.com)

### 5. Expected results

Up to 2500 positive white colonies for a 60 min ligation of 33 ng of supplied control insert (1 kb) and less than 20 negative blue colonies when using cells with a transformation efficiency of  $4 \times 10^7$  cfu/ug.

### 7. Further information

<b>Product Use Limitations</b>	This product is developed, designed and sold exclusively only for research purposes use. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.
<b>Disclaimer</b>	The information provided in this Data Sheet is correct to the best of our knowledge and belief at the date of publication. This information is intended only as a guide and should not be taken as a warranty or quality specification. Canvax Reagents S.L.U. shall not be held liable for any damage resulting from handling or from contact with the above product.

